

Construction of raising and lowering operators for $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$

Pavel ŠTOVÍČEK

Department of Mathematics
Faculty of Nuclear Science, Czech Technical University
Trojanova 13, 120 00 Prague, Czech Republic
E-mail: stovicek@KMLinux.fjfi.cvut.cz

Abstract

In the case of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ at root of unity q -deformed analogues are proposed for the generator of the maximal compact subalgebra, J , and for the raising and lowering operators. We prove an algebraic identity which implies that J has similar properties as in the nondeformed case.

1 Introduction

Despite of an intensive development of the theory of quantum groups during the last decade the results concerning real noncompact forms are rather rare and far away of being complete. Even in the lowest rank cases one encounters serious difficulties (see, for example, references [1, 2]). Naturally, most attention has been paid to groups with a physical interpretation [3, 4]. Here we concentrate on the quantum real group $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$. Even this simple example lacks, to the author's knowledge, a consistent theory on the q -deformed level.

Let us first recall some basic facts concerning the irreducible unitary representations of the classical Lie group $SL(2, \mathbb{R})$ [5, 6]. Let H, E, F be the standard basis elements of $\mathfrak{sl}(2, \mathbb{R})$. All of them are skew self-adjoint with respect to the corresponding $*$ -involution in

$\mathcal{U}(\mathfrak{sl}(2, \mathbb{C}))$. Set

$$J = E - F, \quad W^\pm = H \pm i(E + F).$$

For any irreducible representation ϱ of $SL(2, \mathbb{R})$ in a Hilbert space $\mathcal{H}$ the spectrum of $-i\varrho(J)$ is simple and contained in $\mathbb{Z}$. If $\{v_n\}_{n \in M}$, $M \subset \mathbb{Z}$, is a basis of $\mathcal{H}$ formed by suitably normalised eigenvectors of $-i\varrho(J)$ then

$$-i\varrho(J)v_n = n v_n, \quad \varrho(W^\pm)v_n = (s + 1 \pm n)v_{n \pm 2},$$

where $s \in \mathbb{C}$ is a fixed parameter.

Our main goal here is to propose q -deformed analogues for the generator of the compact subalgebra $\mathfrak{so}(2)$, J , and for the raising and lowering operators $W^\pm$. This short contribution does not aim at all, however, to develop a complete representation theory for $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$; this task is deferred to another publication.

2 $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$ at root of unity

We shall consider the case when q is a root of unity. We assume that

$$q = \exp(2\pi i P/Q)$$

where $Q \in \mathbb{N}$ is odd, $P \in \{1, 2, \dots, Q-1\}$ and P and Q are relatively prime integers. Let K, K^{-1}, E, F be the standard generators of the complex Hopf algebra $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{C}))$. The $*$ -involutions defining the real form is prescribed by the relations

$$K^* = K, \quad E^* = -q^{-1}E, \quad F^* = -qF.$$

Here we prefer another set of generators generating a “slightly smaller” Hopf algebra $\mathcal{U}$, namely

$$X = -i q^{-1} E K^{-1}, \quad Y = -i q F K^{-1}, \quad Z = K^{-2}.$$

Thus the defining relations for $\mathcal{U}$ are

$$\begin{aligned} ZZ^{-1} &= Z^{-1}Z = 1, \quad ZX = q^{-2}XZ, \quad ZY = q^2YZ, \\ q^{-1}XY - qYX &= \frac{1}{q - q^{-1}}(Z^2 - 1), \end{aligned} \tag{1}$$

the comultiplication is defined by

$$\Delta Z = Z \otimes Z, \Delta X = 1 \otimes X + X \otimes Z, \Delta Y = 1 \otimes Y + Y \otimes Z,$$

the antipode is defined by

$$S(Z) = Z^{-1}, S(X) = -XZ^{-1}, S(Y) = -YZ^{-1},$$

and the counit is defined by

$$\varepsilon(Z) = 1, \varepsilon(X) = \varepsilon(Y) = 0.$$

The $*$ -involution defining the real form now reads

$$Z^* = Z, X^* = X, Y^* = Y. \quad (2)$$

The irreducible representations are divided into two families (see the original paper [7] or the monographs [8, 9]). The representations from the first family are labelled by a discrete index $r = 0, 1, \dots, Q-1$, the representation space is $\mathcal{H}_r = \text{span}\{v_0, v_1, \dots, v_r\}$, and the representation is defined by

$$Z \cdot v_j = \pm q^{r-2j} v_j, X \cdot v_j = -q^{r-2j-1} [r-j]_q v_{j+1}, Y \cdot v_j = [j]_q v_{j-1}. \quad (3)$$

The representations from the second family are labelled by 3 complex parameters λ, a, b , the representation space is always $\mathcal{H} = \{v_0, v_1, \dots, v_{Q-1}\}$, the representation is defined by

$$\begin{aligned} Z \cdot v_j &= \lambda q^{2j} v_j, \\ X \cdot v_j &= -i q^{j-1} \left(ab - [j]_q \frac{\lambda q^{j-1} - \lambda^{-1} q^{-j+1}}{q - q^{-1}} \right) v_{j-1} \quad \text{for } j \neq 0, \\ X \cdot v_0 &= -i q^{-1} a v_{Q-1}, \\ Y \cdot v_j &= -i \lambda q^{j+1} v_{j+1} \quad \text{for } j \neq Q-1, \\ Y \cdot v_{Q-1} &= -i \lambda b v_0. \end{aligned} \quad (4)$$

The two families intersect for the values of parameters $r = Q-1$ in (3) and $a = b = 0, \lambda = \pm q^{-Q+1}$ in (4).

Unfortunately, as pointed out in reference [10], the only representations from this list which are unitarizable with respect to the $*$ -involution (2) are one-dimensional. This absolutely unsatisfactory situation of course appeals for a modified definition of $\mathcal{U}_q(\mathfrak{sl}(2, \mathbb{R}))$.

A possible way out has been proposed in reference [10]. The set of generators is augmented by introducing an additional generator T which satisfies

$$T^2 = 1, \Delta T = T \otimes T, S(T) = T, \varepsilon(T) = 1.$$

The augmented Hopf algebra will be called $\mathcal{U}^\natural$. The modified $*$ -involution takes the form

$$X^* = T X T, Y^* = T Y T, Z^* = T Z T, T^* = T. \quad (5)$$

As shown in papers [10, 11], all representations from the first family (3) are unitarizable in this modified sense and may be regarded as q -deformations of representations of $SL(2, \mathbb{R})$ belonging to the discrete series.

3 Raising and lowering operators

Set

$$J = (qX - q^{-1}Y)Z^{-1} = Z^{-1}(q^{-1}X - qY).$$

One easily finds that

$$\Delta J = Z^{-1} \otimes J + J \otimes 1, S(J) = -ZJ, \varepsilon(J) = 1. \quad (6)$$

For the extended algebra $\mathcal{U}^\natural$ it seems to be reasonable to impose the condition

$$JT = TJ,$$

i.e. $J^* = J$.

Note that conversely one can express

$$X = \frac{1}{q^2 - q^{-2}}(qJZ - q^{-1}ZJ), Y = \frac{1}{q^2 - q^{-2}}(q^{-1}JZ - qZJ).$$

It is also straightforward to derive the relations between J and Z . The second and third relations in (1) give

$$Z^2J - (q^2 + q^{-2})ZJZ + JZ^2 = 0 \quad (7)$$

while the last one implies

$$(q^2 + 1 + q^{-2})ZJ^2Z - JZJZ - JZ^2J - ZJZJ = (q + q^{-1})^2(Z^2 - 1). \quad (8)$$

All the above manipulations are reversible and this shows

Proposition 1. *The Hopf algebra $\mathcal{U}$ is isomorphic to the Hopf algebra $\mathcal{U}'$ generated by Z and J with the defining relations (7), (8), (6) and the known relations for Z ($Z Z^{-1} = Z^{-1} Z = 1$, $\Delta Z = Z \otimes Z$, $S(Z) = Z^{-1}$, $\varepsilon(Z) = 1$).*

The quantum numbers $[x]_q = (q^x - q^{-x})/(q - q^{-1})$ are supposed to be defined for any $x \in \mathbb{C}$ for we set, by definition, $q^x = \exp(2\pi i(P/Q)x)$. The main result of this contribution is the following identity.

Proposition 2. *For all $x \in \mathbb{C}$ it holds true that*

$$\begin{aligned} & Z(J - [x + 2]_q)(J - [x]_q)(J - [x - 2]_q)Z \\ &= \left((J - [x]_q)Z(J - [x]_q)Z - [2]_q^2 \right)(J - [x]_q), \end{aligned} \quad (9)$$

Proof. To prove (9) we expend the both sides in the variable $y = q^x$,

$$\text{LHS} - \text{RHS} = c_{-2}y^{-2} + c_{-1}y^{-1} + c_0 + c_1y + c_2y^2,$$

and we find that

$$\begin{aligned} (q - q^{-1})^2 c_{-2} &= (q - q^{-1})^2 c_2 = -JZ^2 + (q^2 + q^{-2})ZJZ - Z^2J, \\ (q - q^{-1})c_{-1} &= -(q - q^{-1})c_1 \\ &= (q^2 + 1 + q^{-2})ZJ^2Z - JZJZ - JZ^2J - ZJZJ - [2]_q^2(Z^2 - 1), \end{aligned}$$

and

$$\begin{aligned} (q - q^{-1})^2 c_0 &= 2(JZ^2 - (q^2 + q^{-2})ZJZ + Z^2J) \\ &\quad + (q - q^{-1})^2 (ZJ^3Z - [2]_q^2 ZJZ - JZJZJ + [2]_q^2 J). \end{aligned}$$

Thus the coefficients $c_{\pm 2}$, $c_{\pm 1}$ vanish owing to (7) and (8), the coefficient c_0 vanishes if it is true in the particular case $x = 0$. The following lemma concludes the proof. $\square$

Lemma 3. *It holds true that*

$$Z(J^2 - (q + q^{-1})^2)JZ = (JZJZ - (q + q^{-1})^2)J.$$

Proof. To prove the particular case of (9) with $x = 0$ set

$$V = ZJ^3Z - [2]_q^2 ZJZ + [2]_q^2 J - JZJZJ.$$

Then it holds

$$\begin{aligned} ZV + VZ = & ((q^2 + 1 + q^{-2}) ZJ^2Z - JZ^2J - ZJZJ - JZJZ + (q + q^{-1})^2(1 - Z^2)) \\ & \times JZ + ZJ \\ & \times ((q^2 + 1 + q^{-2}) ZJ^2 - JZ^2J - ZJZJ - JZJZ + (q + q^{-1})^2(1 - Z^2)) \\ & + (JZ^2 - (q^2 + q^{-2})ZJZ + Z^2J)J^2Z \\ & + ZJ^2(JZ^2 - (q^2 + q^{-2})ZJZ + Z^2J) \end{aligned}$$

and thus $ZV + VZ = 0$. This clearly implies

$$Z^k V = (-1)^k VZ, \quad \text{for all } k \in \mathbb{Z}.$$

In particular, $Z^Q V = -VZ^Q$. But Z^Q belongs to the centre of the algebra $\mathcal{U}$ and therefore $Z^Q V = VZ^Q$. Hence $Z^Q V = 0$ and consequently $V = 0$. $\square$

Some straightforward consequences follow from identity (9). Consider an irreducible representation of $\mathcal{U}$ whose dimension equals d . If v is an eigenvector of J with an eigenvalue $[x]_q$ then the vectors

$$(J - [x]_q)(J - [x - 2]_q)Z \cdot v, \quad (J - [x]_q)(J - [x + 2]_q)Z \cdot v, \quad (10)$$

either vanish or are eigenvectors of J corresponding to the eigenvalues $[x + 2]_q$ and $[x - 2]_q$, respectively. The spectrum of J is of the form

$$[x]_q, [x + 2]_q, \dots, [x + 2d - 2]_q,$$

and hence the operator J has similar properties (up to the factor i) as its nondeformed counterpart. For example, in the case of the representations from the first family (3) the operator J is known to have the spectrum (10) with $x = Q - d + 1$ [10]. Furthermore, the operators $(J - [x]_q)(J - [x - 2]_q)Z$ and $(J - [x]_q)(J - [x + 2]_q)Z$ play the role of raising and lowering operators, respectively. Finally, the matrix Z , if expressed in the corresponding eigenbasis of J , is tridiagonal in the case of the first family of representations (3) or tridiagonal with the cyclic convention modulo Q in the case of the second family (4).

Acknowledgements

The research was partially supported by Grant GACR 201/01/01308.

References

- [1] S.L. Woronowicz: Lett. Math. Phys. **23** (1991) 251.
- [2] L.I. Korogodsky: Commun. Math. Phys. **163** (1994) 433.
- [3] V.K. Dobrev, P. Moylan: Phys. Lett. B **315** (1993) 292.
- [4] V.K. Dobrev and R. Floreanini: J. Phys. **A27** (1994) 4831.
- [5] V. Bargmann: Ann. Math. **48** (1947) 568.
- [6] S. Lang: $SL_2(\mathbf{R})$. Addison-Wesley, Reading, 1975.
- [7] P. Roche, D. Arnaudon: Lett. Math. Phys. **17** (1989) 295.
- [8] V. Chari, A. Pressley: *A Guide to Quantum Groups*. Cambridge University Press, Cambridge, 1994.
- [9] A. Klimyk, K. Schmüdgen: *Quantum Groups and Their Representations*. Springer-Verlag, Berlin Heidelberg, 1997.
- [10] P. Šťovíček: Czech. J. Phys. **50** (2000) 1353.
- [11] P. Šťovíček: in *Proc. Internat. Sympos. Quantum Theory and Symmetries*, Goslar 1999 (Eds. H.-D. Doebner et al.), World Scientific, Singapore, 2000, p.470.